VOORHEES MIDDLE SCHOOL HOLLY OAK DRIVE, VOORHEES, NJ 08043 Athletic Team Parent/Guardian Permission Form

Name o	of Student		Telephone				
Date of	Birth	eroom Teacher					
Address							
	FIONS FOR ATHLETIC PRE-PA s to participate in interscholastic		The following forms must be completed in order for ms.				
	rticipation Physical Evaluation the Physician for review at time		ust be completed and signed by Parent/Guardian and				
* Physical Examination Form and Clearance Form: These 2 pages must be completed by the examining licensed provide MD, DO, APN, or PA. This physical exam must have been completed WITHIN 365 days BEFORE the first day of try-outs							
*Concu	ssion Acknowledgement Form	& Sudden Cardiac Death Pa	mphlet sign-off Sheet: Signed by parent & student				
	H HISTORY UPDATE QUESTIO Update must be submitted 90		for additional sports provided a current physical is of each sport.				
To be 0	Completed by Parent/Guardian						
This is to	certify that we, the undersigned, ha	ve given orhees Interscholastic/Intramural t	(name) permission to playeam.				
	arent/guardian we realize there are c ibility for our child's safety.	ertain physical hazards connected	with this activity and are willing to assume absolutely all				
Date	Parent/G	uardian Signature					
To be 0	Completed by Student						
2. 3. 4.	I shall not neglect my scholastic wor I shall, as a member of the group or I shall report regularly to all schedule I shall be responsible for all athletic I shall attempt to improve the team of	squad, abide by training regulation ed practices and contests. supplies issued to me and shall re					
Date	Student's	s Signature					
	Fall Sports Boys/Girls Soccer Field Hockey	Winter Sports Boys/Girls Basketball Cheerleading	Spring Sports Boys/Girls Track Baseball				

Wrestling

Softball

Voorhees Middle School

Sports-Related Concussion and Head Injury Fact Sheet and Parent/Guardian Acknowledgement Form

A concussion is a brain injury that can be caused by a blow to the head or body that disrupts normal functioning of the brain. Concussions are a type of Traumatic Brain Injury (TBI), which can range from mild to severe and can disrupt the way the brain normally functions. Concussions can cause significant and sustained neuropsychological impairment affecting problem solving, planning, memory, attention, concentration, and behavior.

The Centers for Disease Control and Prevention estimates that 300,000 concussions are sustained during sports related activities nationwide, and more than 62,000 concussions are sustained each year in high school contact sports. Second-impact syndrome occurs when a person sustains a second concussion while still experiencing symptoms of a previous concussion. It can lead to severe impairment and even death of the victim.

Legislation (P.L. 2010, Chapter 94) signed on December 7, 2010, mandated measures to be taken in order to ensure the safety of K-12 student-athletes involved in interscholastic sports in New Jersey. It is imperative that athletes, coaches, and parent/guardians are educated about the nature and treatment of sports related concussions and other head injuries. The legislation states that:

- All Coaches, Athletic Trainers, School Nurses, and School/Team Physicians shall complete an Interscholastic Head Injury Safety Training Program by the 2011-2012 school year.
- All school districts, charter, and non-public schools that participate in interscholastic sports will distribute annually this educational fact to all student athletes and obtain a signed acknowledgement from each parent/guardian and student-athlete.
- Each school district, charter, and non-public school shall develop a written policy describing the prevention and treatment of sports-related concussion and other head injuries sustained by interscholastic student-athletes.
- Any student-athlete who participates in an interscholastic sports program and is suspected of sustaining a
 concussion will be immediately removed from competition or practice. The student-athlete will not be
 allowed to return to competition or practice until he/she has written clearance from a physician trained in
 concussion treatment and has completed his/her district's graduated return-to-play protocol.

Quick Facts

- Most concussions do not involve loss of consciousness
- You can sustain a concussion even if you do not hit your head
- A blow elsewhere on the body can transmit an "impulsive" force to the brain and cause a concussion

Signs of Concussions (Observed by Coach, Athletic Trainer, Parent/Guardian)

- Appears dazed or stunned
- Forgets plays or demonstrates short term memory difficulties (e.g. unsure of game, opponent)
- Exhibits difficulties with balance, coordination, concentration, and attention
- Answers questions slowly or inaccurately
- Demonstrates behavior or personality changes
- Is unable to recall events prior to or after the hit or fall

Symptoms of Concussion (Reported by Student-Athlete)

- Headache
- Nausea/vomiting
- Balance problems or dizziness
- Double vision or changes in vision

- Sensitivity to light/sound
- Feeling of sluggishness or fogginess
- Difficulty with concentration, short term memory, and/or confusion

Voorhees Middle School

What Should a Student-Athlete do if they think they have a concussion?

- Don't hide it. Tell your Athletic Trainer, Coach, School Nurse, or Parent/Guardian.
- **Report it**. Don't return to competition or practice with symptoms of a concussion or head injury. The sooner you report it, the sooner you may return-to-play.
- Take time to recover. If you have a concussion your brain needs time to heal. While your brain is healing you are much more likely to sustain a second concussion. Repeat concussions can cause permanent brain injury.

What can happen if a student-athlete continues to play with a concussion or returns to play to soon?

- Continuing to play with the signs and symptoms of a concussion leaves the student-athlete vulnerable to second impact syndrome.
- Second impact syndrome is when a student-athlete sustains a second concussion while still having symptoms from a previous concussion or head injury.
- Second impact syndrome can lead to severe impairment and even death in extreme cases.

Should there be any temporary academic accommodations made for Student-Athletes who have suffered a concussion?

- To recover cognitive rest is just as important as physical rest. Reading, texting, testing-even watching movies can slow down a student-athletes recovery.
- Stay home from school with minimal mental and social stimulation until all symptoms have resolved.
- Students may need to take rest breaks, spend fewer hours at school, be given extra time to complete assignments, as well as being offered other instructional strategies and classroom accommodations.

Student-Athletes who have sustained a concussion should complete a graduated return-to-play before they may resume competition or practice, according to the following protocol:

- **Step 1**: Completion of a full day of normal cognitive activities (school day, studying for tests, watching practice, interacting with peers) without reemergence of any signs or symptoms. If no return of symptoms, next day advance.
- **Step 2:** Light Aerobic exercise, which includes walking, swimming, and stationary cycling, keeping the intensity below 70% maximum heart rate. No resistance training. The objective of this step is increased heart rate.
- **Step 3:** Sport-specific exercise including skating, and/or running: no head impact activities. The objective of this step is to add movement.
- Step 4: Non contact training drills (e.g. passing drills). Student-athlete may initiate resistance training.
- **Step 5:** Following medical clearance (consultation between school health care personnel and student-athlete's physician), participation in normal training activities. The objective of this step is to restore confidence and assess functional skills by coaching and medical staff.
- **Step 6:** Return to play involving normal exertion or game activity.

www.cdc.gov/concussion/sports/index.html		www.nfhs.com	
www.ncaa.org/health-safety	www.bianj.org www.atsnj.org		
Signature of Student-Athlete	Print Student-A	Student-Athlete's Name	
Signature of Parent/Guardian	Print Parent/Gu	ordian's Nama	Date

Website Resources

- Sudden Death in Athletes http://tinyurl.com/m2gjmvq
- Hypertrophic Cardiomyopathy Association www.4hcm.org
- American Heart Association www.heart.org

Collaborating Agencies:

American Academy of Pediatrics New Jersey Chapter

3836 Quakerbridge Road, Suite 108 Hamilton, NJ 08619 (p) 609-842-0014 (f) 609-842-0015 www.aapnj.org



American Heart Association

1 Union Street, Suite 301 Robbinsville, NJ, 08691 (p) 609-208-0020 www.heart.org



New Jersey Department of Education

PO Box 500 Trenton, NJ 08625-0500 (p) 609-292-5935 www.state.nj.us/education/



New Jersey Department of Health

P. O. Box 360 Trenton, NJ 08625-0360 (p) 609-292-7837 www.state.nj.us/health




Lead Author: American Academy of Pediatrics, New Jersey Chapter

Written by: Initial draft by Sushma Raman Hebbar, MD & Stephen G. Rice, MD PhD

Additional Reviewers: NJ Department of Education, NJ Department of Health and Senior Services, American Heart Association/New Jersey Chapter, NJ Academy of Family Practice, Pediatric Cardiologists, New Jersey State School Nurses

Revised 2014: Nancy Curry, EdM; Christene DeWitt-Parker, MSN, CSN, RN; Lakota Kruse, MD, MPH; Susan Martz, EdM; Stephen G. Rice, MD; Jeffrey Rosenberg, MD, Louis Teichholz, MD; Perry Weinstock, MD




American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®







Sudden death in young athletes between the ages of 10 and 19 is very rare. What, if anything, can be done to prevent this kind of tragedy?

What is sudden cardiac death in the young athlete?

Sudden cardiac death is the result of an unexpected failure of proper heart function, usually (about 60% of the time) during or immediately after exercise without trauma. Since the heart stops pumping adequately, the athlete quickly collapses, loses consciousness, and ultimately dies unless normal heart rhythm is restored using an automated external defibrillator (AED).

How common is sudden death in young athletes?

Sudden cardiac death in young athletes is very rare. About 100 such deaths are reported in the United States per year. The chance of sudden death occurring to any individual high school athlete is about one in 200,000 per year.

Sudden cardiac death is more common: in males than in females; in football and basketball than in other sports; and in African-Americans than in other races and ethnic groups.

What are the most common causes?

Research suggests that the main cause is a loss of proper heart rhythm, causing the heart to quiver instead of pumping blood to the brain and body. This is called ventricular fibrillation (ven-TRICK-you-lar fibroo-LAY-shun). The problem is usually caused by one of several cardiovascular abnormalities and electrical diseases of the heart that go unnoticed in healthy-appearing athletes.

The most common cause of sudden death in an athlete is hypertrophic cardiomyopathy (hi-per-TRO-fic CAR- dee-oh-my-OP-a-thee) also called HCM. HCM is a disease of the heart, with abnormal thickening of the heart muscle, which can cause serious heart rhythm problems and blockages to blood flow. This genetic disease runs in families and usually develops gradually over many years.

The second most likely cause is congenital (con-JEN-it-al) (i.e., present from birth) abnormalities of the coronary

arteries. This means that these blood vessels are connected to the main blood vessel of the heart in an abnormal way. This differs from blockages that may occur when people get older (commonly called "coronary artery disease," which may lead to a heart attack).

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES

Other diseases of the heart that can lead to sudden death in young people include:

- Myocarditis (my-oh-car-DIE-tis), an acute inflammation of the heart muscle (usually due to a virus).
- Dilated cardiomyopathy, an enlargement of the heart for unknown reasons.
- Long QT syndrome and other electrical abnormalities of the heart which cause abnormal fast heart rhythms that can also run in families.
- Marfan syndrome, an inherited disorder that affects heart valves, walls of major arteries, eyes and the skeleton. It is generally seen in unusually tall athletes, especially if being tall is not common in other family members.

Are there warning signs to watch for?

In more than a third of these sudden cardiac deaths, there were warning signs that were not reported or taken seriously. Warning signs are:

- Fainting, a seizure or convulsions during physical activity;
- Fainting or a seizure from emotional excitement, emotional distress or being startled;
- Dizziness or lightheadedness, especially during exertion;
- Chest pains, at rest or during exertion;
- Palpitations awareness of the heart beating unusually (skipping, irregular or extra beats) during athletics or during cool down periods after athletic participation;
- \bullet Fatigue or tiring more quickly than peers; or
- Being unable to keep up with friends due to shortness of breath (labored breathing).

What are the current recommendations for screening young athletes?

New Jersey requires all school athletes to be examined by their primary care physician ("medical home") or school physician at least once per year. The New Jersey Department of Education requires use of the specific Preparticipation Physical Examination Form (PPE).

This process begins with the parents and student-athletes answering questions about symptoms during exercise (such as chest pain, dizziness, fainting, palpitations or shortness of breath); and questions about family health history.

The primary healthcare provider needs to know if any family member died suddenly during physical activity or during a seizure. They also need to know if anyone in the family under the age of 50 had an unexplained sudden death such as drowning or car accidents. This information must be provided annually for each exam because it is so essential to identify those at risk for sudden cardiac death.

The required physical exam includes measurement of blood pressure and a careful listening examination of the heart, especially for murmurs and rhythm abnormalities. If there are no warning signs reported on the health history and no abnormalities discovered on exam, no further evaluation or testing is recommended.

Are there options privately available to screen for cardiac conditions?

Technology-based screening programs including a 12-lead electrocardiogram (ECG) and echocardiogram (ECHO) are noninvasive and painless options parents may consider in addition to the required

PPE. However, these procedures may be expensive and are not currently advised by the American Academy of Pediatrics and the American College of Cardiology unless the PPE reveals an indication for these tests. In addition to the expense, other limitations of technology-based tests include the possibility of "false positives" which leads to unnecessary stress for the student and parent or guardian as well as unnecessary restriction from athletic participation.

The United States Department of Health and Human Services offers risk assessment options under the Surgeon General's Family History Initiative available at http://www.hhs.gov/familyhistory/index.html.

When should a student athlete see a heart specialist?

If the primary healthcare provider or school physician has concerns, a referral to a child heart specialist, a pediatric cardiologist, is recommended. This specialist will perform a more thorough evaluation, including an electrocardiogram (ECG), which is a graph of the electrical activity of the heart. An echocardiogram, which is an ultrasound test to allow for direct visualization of the heart structure, will likely also be done. The specialist may also order a treadmill exercise test and a monitor to enable a longer recording of the heart rhythm. None of the testing is invasive or uncomfortable.

Can sudden cardiac death be prevented just through proper screening?

A proper evaluation should find most, but not all, conditions that would cause sudden death in the athlete. This is because some diseases are difficult to uncover and may only develop later in life. Others can develop following a normal screening evaluation, such as an infection of the heart muscle from a virus.

This is why screening evaluations and a review of the family health history need to be performed on a yearly basis by the athlete's primary healthcare provider. With proper screening and evaluation, most cases can be identified and prevented.

Why have an AED on site during sporting events?

The only effective treatment for ventricular fibrillation is immediate use of an automated external defibrillator (AED). An AED can restore the heart back into a normal rhythm. An AED is also life-saving for ventricular fibrillation caused by a blow to the chest over the heart (commotio cordis).

N.J.S.A. 18A:40-41a through c, known as "Janet's Law," requires that at any schoolsponsored athletic event or team practice in New Jersey public and nonpublic schools including any of grades K through 12, the following must be available:

- An AED in an unlocked location on school property within a reasonable proximity to the athletic field or gymnasium; and
- A team coach, licensed athletic trainer, or other designated staff member if there is no coach or licensed athletic trainer present, certified in cardiopulmonary resuscitation (CPR) and the use of the AED; or
- A State-certified emergency services provider or other certified first responder.

The American Academy of Pediatrics recommends the AED should be placed in central location that is accessible and ideally no more than a 1 to 1½ minute walk from any location and that a call is made to activate 911 emergency system while the AED is being retrieved.

State of New Jersey DEPARTMENT OF EDUCATION

$\frac{\textbf{Sudden Cardiac Death Pamphlet}}{\textbf{Sign-Off Sheet}}$

Name of School District:
Name of Local School:
I/We acknowledge that we received and reviewed the Sudden Cardiac Death in Young Athletes pamphlet.
Student Signature:
Parent or Guardian Signature:
Date:

SPORTS-RELATED

EYE INJURIES:

AN EDUCATIONAL FACT SHEET FOR PARENTS



Participating in sports and recreational activities is an important part of a healthy, physically active lifestyle for children. Unfortunately, injuries can, and do, occur. Children are at particular risk for sustaining a sports-related eye injury and most of these injuries can be prevented. Every year, more than 30,000 children sustain serious sports-related eye injuries. Every 13 minutes, an emergency room in the United States treats a sports-related eye injury. According to the National Eye Institute, the sports with the highest rate of eye injuries are: baseball/softball, ice hockey, racquet sports, and basketball, followed by fencing, lacrosse, paintball and boxing.

Thankfully, there are steps that parents can take to ensure their children's safety on the field, the court, or wherever they play or participate in sports and recreational activities.

Prevention of Sports-Related Eye Injuries Approximately 90% of sports-related eye injuries can be prevented with simple precautions, such as using protective eyewear.² Each sport has a certain type of recommended protective eyewear, as determined by the American Society for Testing and Materials (ASTM). Protective eyewear should sit comfortably on the face. Poorly fitted equipment may be uncomfortable, and may not offer the best eye protection. Protective eyewear for sports includes, among other things, safety goggles and eye guards, and it should be made of polycarbonate lenses, a strong, shatterproof plastic. Polycarbonate lenses are much stronger than regular lenses.³

Health care providers (HCP), including family physicians, ophthalmologists, optometrists, and others, play a critical role in advising students, parents and guardians about the proper use of protective eyewear. To find out what kind of eye protection is recommended, and permitted for your child's sport, visit the National Eye Institute at http://www.nei.nih.gov/sports/findingprotection.asp. Prevent Blindness America also offers tips for choosing and buying protective eyewear at http://www.preventblindness.org/tips-buying-sports-eye-protectors, and http://www.preventblindness.org/ recommended-sports-eye-protectors.

It is recommended that all children participating in school sports or recreational sports wear protective eyewear. Parents and coaches need to make sure young athletes protect their eyes, and properly gear up for the game. Protective eyewear should be part of any uniform to help reduce the occurrence of sports-related eye injuries. Since many youth teams do not require eye protection, parents may need to ensure that their children wear safety glasses or goggles whenever they play sports. Parents can set a good example by wearing protective eyewear when they play sports.

¹ National Eye Institute, National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nei.nih.gov/sports/pdf/sportsrelatedeyeInjuries.pdf, December 26, 2013.

² Rodriguez, Jorge O., D.O., and Lavina, Adrian M., M.D., Prevention and Treatment of Common Eye Injuries in Sports, http://www.aafp.org/afp/2003/0401/p1481.html, September 4, 2014; National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nei.nih.gov/sports/pdf/sportsrelatedeyeInjuries.pdf, December 26, 2013.

Bedinghaus, Troy, O.D., Sports Eye Injuries, http://vision.about.com/od/emergencyeyecare/a/Sports_Injuries.htm, December 27, 2013.

The most common types of eye injuries that can result from sports injuries are

blunt injuries, corneal abrasions and penetrating injuries.



- ♦ Blunt injuries: Blunt injuries occur when the eye is suddenly compressed by impact from an object. Blunt injuries, often caused by tennis balls, racquets, fists or elbows, sometimes cause a black eye or hyphema (bleeding in front of the eye). More serious blunt injuries often break bones near the eye, and may sometimes seriously damage important eye structures and/or lead to vision loss.
- ◆ Corneal abrasions: Corneal abrasions are painful scrapes on the outside of the eye, or the cornea. Most corneal abrasions eventually heal on their

own, but a doctor can best assess the extent of the abrasion, and may prescribe medication to help control the pain. The most common cause of a sports-related corneal abrasion is being poked in the eye by a finger.

- ◆ Penetrating injuries: Penetrating injuries are caused by a foreign object piercing the eye. Penetrating injuries are very serious, and often result in severe damage to the eye. These injuries often occur when eyeglasses break while they are being worn. Penetrating injuries must be treated quickly in order to preserve vision.⁴
- Pain when looking up and/or down, or difficulty seeing;
- Tenderness;
- Sunken eye;
- Double vision:
- Severe eyelid and facial swelling;
- Difficulty tracking;

Signs or Symptoms of an Eye Injury



- The eye has an unusual pupil size or shape;
- Blood in the clear part of the eye;
- Numbness of the upper cheek and gum; and/or
- Severe redness around the white part of the eye.

What to do if a Sports-Related Eye Injury Occurs

If a child sustains an eye injury, it is recommended that he/she receive immediate treatment from a licensed HCP (e.g., eye doctor) to reduce the risk of serious damage, including blindness. It is also recommended that the child, along with his/her parent or guardian, seek guidance from the HCP regarding the appropriate amount of time to wait before returning to sports competition or practice after sustaining an eye injury. The school nurse and the child's teachers should also be notified when a child sustains an eye injury. A parent or guardian should also provide the school nurse with a physician's note detailing the nature of the eye injury, any diagnosis, medical orders for

the return to school, as well as any prescription(s) and/or treatment(s) necessary to promote healing, and the safe resumption of normal activities, including sports and recreational activities.

Return to Play and Sports

According to the American Family Physician Journal, there are several guidelines that should be followed when students return to play after sustaining an eye injury. For

example, students who have sustained significant ocular injury should receive a full examination and clearance by an ophthalmologist or optometrist. In addition, students should not return to play until the period of time recommended by their HCP has elapsed. For more minor eye injuries, the athletic trainer may determine that

it is safe for a student to resume play based on the nature of the injury, and how the student feels. No matter what degree of eye injury is sustained, it is recommended that students wear protective eyewear when returning to play and immediately report any concerns with their vision to their coach and/or the athletic trainer.

Additional information on eye safety can be found at http://isee.nei.nih.gov and http://www.nei.nih.gov/sports.

SCHOLASTIC STUDENT-ATHLETE SAFETY ACT INFORMATION FACT SHEET FOR PARENTS/GUARDIANS

Prior to participation on a school-sponsored interscholastic or intramural athletic team or squad, each student-athlete in grades six through 12 must present a completed pre-participation physical evaluation (PPE) form to the designated school staff member. Important information regarding the PPE is provided below, and you should <u>feel free to share with your child's medical home health care provider</u>.

- 1. The PPE may ONLY be completed by a licensed physician, advanced practice nurse (APN) or physician assistant (PA) that has completed the Student-Athlete Cardiac Assessment professional development module. It is recommended that you verify that your medical provider has completed this module before scheduling an appointment for a PPE.
- 2. The required PPE must be conducted within 365 days prior to the first official practice in an athletic season. The PPE form is available in English and Spanish at http://www.state.nj.us/education/students/safety/health/records/athleticphysicalsform.pdf.
- 3. The parent/guardian must complete the *History Form* (page one), and insert the date of the required physical examination at the top of the page.
- 4. The parent/guardian must complete *The Athlete with Special Needs: Supplemental History Form* (page two), if applicable, for a student with a disability that limits major life activities, and insert the date of the required physical examination on the top of the page.
- 5. The licensed physician, APN or PA who performs the physical examination must complete the remaining two pages of the PPE, and insert the date of the examination on the *Physical Examination Form* (page three) and *Clearance Form* (page four).
- 6. The school district must provide written notification to the parent/guardian, signed by the school physician, indicating approval of the student's participation in a school-sponsored interscholastic or intramural athletic team or squad based upon review of the medical report, or must provide the reason(s) for the disapproval of the student's participation.
- 7. For student-athletes that had a medical examination completed more than 90 days prior to the first official practice in an athletic season, the *Health History Update Questionnaire* (HHQ) form must be completed, and signed by the student's parent/guardian. The HHQ must be reviewed by the school nurse and, if applicable, the school's athletic trainer. The HHQ is available at http://www.state.nj.us/education/students/safety/health/records/HealthHistoryUpdate.pdf.

For more information, please review the *Frequently Asked Questions* which are available at http://www.state.nj.us/education/students/safety/health/services/athlete/faq.pdf.

ATTENTION PARENT/GUARDIAN: The preparticipation physical examination (page 3) must be completed by a health care provider who has completed the Student-Athlete Cardiac Assessment Professional Development Module.

■ PREPARTICIPATION PHYSICAL EVALUATION HISTORY FORM

ame				Date of birth		
ex Age Grade	Schoo	ol		Sport(s)		
Medicines and Allergies: Please list all of the prescr	iption and over-th	ne-cou	ınter m	edicines and supplements (herbal and nutritional) that you are currently	taking	
Do you have any allergies? ☐ Yes ☐ No ☐ If y☐ Medicines ☐ Polle		fy spe		ergy below. ☐ Food ☐ Stinging Insects		
plain "Yes" answers below. Circle questions you don	't know the answ	vers to).			
ENERAL QUESTIONS		Yes	No	MEDICAL QUESTIONS	Yes	
Has a doctor ever denied or restricted your participation in any reason?	sports for			26. Do you cough, wheeze, or have difficulty breathing during or after exercise?		L
2. Do you have any ongoing medical conditions? If so, pleas				27. Have you ever used an inhaler or taken asthma medicine?		╄
below: ☐ Asthma ☐ Anemia ☐ Diabetes ☐ In Other:	nfections			28. Is there anyone in your family who has asthma?		╀
3. Have you ever spent the night in the hospital?				29. Were you born without or are you missing a kidney, an eye, a testicle (males), your spleen, or any other organ?		
1. Have you ever had surgery?				30. Do you have groin pain or a painful bulge or hernia in the groin area?		Τ
EART HEALTH QUESTIONS ABOUT YOU		Yes	No	31. Have you had infectious mononucleosis (mono) within the last month?		
5. Have you ever passed out or nearly passed out DURING or				32. Do you have any rashes, pressure sores, or other skin problems?		I
AFTER exercise?	in your			33. Have you had a herpes or MRSA skin infection?		
6. Have you ever had discomfort, pain, tightness, or pressure chest during exercise?	e iii your			34. Have you ever had a head injury or concussion?		1
7. Does your heart ever race or skip beats (irregular beats) of	luring exercise?			35. Have you ever had a hit or blow to the head that caused confusion, prolonged headache, or memory problems?		
3. Has a doctor ever told you that you have any heart proble	ms? If so,			36. Do you have a history of seizure disorder?		†
check all that apply: ☐ High blood pressure ☐ A heart murmur				37. Do you have headaches with exercise?		T
☐ High cholesterol ☐ A heart infection ☐ Kawasaki disease Other:				38. Have you ever had numbness, tingling, or weakness in your arms or legs after being hit or falling?		
Has a doctor ever ordered a test for your heart? (For exame chocardiogram)	nple, ECG/EKG,			39. Have you ever been unable to move your arms or legs after being hit or falling?		
Do you get lightheaded or feel more short of breath than a during exercise?	expected			40. Have you ever become ill while exercising in the heat?		╀
Have you ever had an unexplained seizure?				41. Do you get frequent muscle cramps when exercising? 42. Do you or someone in your family have sickle cell trait or disease?		+
Do you get more tired or short of breath more quickly that	n your friends			43. Have you had any problems with your eyes or vision?		+
during exercise?	,			44. Have you had any eye injuries?		t
EART HEALTH QUESTIONS ABOUT YOUR FAMILY		Yes	No	45. Do you wear glasses or contact lenses?		t
Has any family member or relative died of heart problems unexpected or unexplained sudden death before age 50 (46. Do you wear protective eyewear, such as goggles or a face shield?		T
drowning, unexplained car accident, or sudden infant dea				47. Do you worry about your weight?		
4. Does anyone in your family have hypertrophic cardiomyor				48. Are you trying to or has anyone recommended that you gain or		
syndrome, arrhythmogenic right ventricular cardiomyopat syndrome, short QT syndrome, Brugada syndrome, or cat				lose weight? 49. Are you on a special diet or do you avoid certain types of foods?		+
polymorphic ventricular tachycardia?				50. Have you ever had an eating disorder?		+
5. Does anyone in your family have a heart problem, pacema	aker, or			51. Do you have any concerns that you would like to discuss with a doctor?		+
implanted defibrillator? 6. Has anyone in your family had unexplained fainting, unexpla	nlained			FEMALES ONLY		
seizures, or near drowning?	Jameu			52. Have you ever had a menstrual period?		T
ONE AND JOINT QUESTIONS		Yes	No	53. How old were you when you had your first menstrual period?		
7. Have you ever had an injury to a bone, muscle, ligament, that caused you to miss a practice or a game?	or tendon			54. How many periods have you had in the last 12 months?		
Have you ever had any broken or fractured bones or dislo	cated joints?			Explain "yes" answers here		
9. Have you ever had an injury that required x-rays, MRI, CT	-					
injections, therapy, a brace, a cast, or crutches?						
Have you ever had a stress fracture? Here you ever heap told that you have at have you had as	v rou for mc-1:					
 Have you ever been told that you have or have you had ar instability or atlantoaxial instability? (Down syndrome or c 						
2. Do you regularly use a brace, orthotics, or other assistive						
3. Do you have a bone, muscle, or joint injury that bothers yo	ou?					
24. Do any of your joints become painful, swollen, feel warm, 25. Do you have any history of juvenile arthritis or connective						_

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

■ PREPARTICIPATION PHYSICAL EVALUATION THE ATHLETE WITH SPECIAL NEEDS: SUPPLEMENTAL HISTORYFORM

Date of Exam					
Name			Date of birth		
Sav Ana	Grade	School			
Sex Age	uraue	301001	Sport(s)		
1. Type of disability					
2. Date of disability					
3. Classification (if avail	lable)				
4. Cause of disability (b	irth, disease, accident/trauma, other)				
5. List the sports you ar	re interested in playing				
				Yes	No
6. Do you regularly use	a brace, assistive device, or prostheti	0?			
7. Do you use any spec	ial brace or assistive device for sports	?			
8. Do you have any rash	hes, pressure sores, or any other skin	problems?			
	ng loss? Do you use a hearing aid?				
10. Do you have a visual					
	ial devices for bowel or bladder functi	on?			
	or discomfort when urinating?				
13. Have you had autono					
		nermia) or cold-related (hypothermia) illnes	ss?		
15. Do you have muscle					
16. Do you have frequen	t seizures that cannot be controlled by	medication?			
Explain "yes" answers h	ere				
Please indicate if you ha	ve ever had any of the following.				
				Yes	No
Atlantoaxial instability					
X-ray evaluation for atlan	toaxial instability				
Dislocated joints (more th	nan one)				
Easy bleeding					
Enlarged spleen					
Hepatitis					
Osteopenia or osteoporos	sis				
Difficulty controlling bow	el				
Difficulty controlling blad					
Numbness or tingling in a					
Numbness or tingling in I	<u> </u>				
Weakness in arms or han	nds				
Weakness in legs or feet					
Recent change in coordin					
Recent change in ability t	to walk				
Spina bifida					
Latex allergy					
Explain "yes" answers h	ere				
I hereby state that, to the	e best of my knowledge, my answer	rs to the above questions are complete a	and correct.		
Signature of athlete		Signature of parent/guardian		Date	

NOTE: The preparticiaption physical examination must be conducted by a health care provider who 1) is a licensed physician, advanced practice nurse, or physician assistant; and 2) completed the Student-Athlete Cardiac Assessment Professional Development Module.

PREPARTICIPATION PHYSICAL EVALUATION


PHYSICAL EXAMINATION FORM Name Date of birth **PHYSICIAN REMINDERS** 1. Consider additional questions on more sensitive issues Do you feel stressed out or under a lot of pressure? Do you ever feel sad, hopeless, depressed, or anxious? • Do you feel safe at your home or residence? • Have you ever tried cigarettes, chewing tobacco, snuff, or dip? • During the past 30 days, did you use chewing tobacco, snuff, or dip? Do you drink alcohol or use any other drugs? • Have you ever taken anabolic steroids or used any other performance supplement? • Have you ever taken any supplements to help you gain or lose weight or improve your performance? • Do you wear a seat belt, use a helmet, and use condoms? 2. Consider reviewing questions on cardiovascular symptoms (questions 5-14). **EXAMINATION** Height Weight □ Male □ Female BP Pulse Vision R 20/ L 20/ Corrected D Y \square N MEDICAL NORMAL ABNORMAL FINDINGS · Marfan stigmata (kyphoscoliosis, high-arched palate, pectus excavatum, arachnodactyly, arm span > height, hyperlaxity, myopia, MVP, aortic insufficiency) Eyes/ears/nose/throat · Pupils equal Hearing Lymph nodes Heart^a Murmurs (auscultation standing, supine, +/- Valsalva) Location of point of maximal impulse (PMI) Pulses · Simultaneous femoral and radial pulses Lungs Abdomen Genitourinary (males only)b . HSV, lesions suggestive of MRSA, tinea corporis Neurologic o MUSCULOSKELETAL Neck Back Shoulder/arm Elbow/forearm Wrist/hand/fingers Hip/thigh Knee Leg/ankle Foot/toes **Functional** · Duck-walk, single leg hop ^aConsider ECG, echocardiogram, and referral to cardiology for abnormal cardiac history or exam. ^bConsider GU exam if in private setting. Having third party present is recommended.
^cConsider cognitive evaluation or baseline neuropsychiatric testing if a history of significant concussion. ☐ Cleared for all sports without restriction ☐ Cleared for all sports without restriction with recommendations for further evaluation or treatment for _ □ Not cleared □ Pending further evaluation □ For any sports □ For certain sports _ Recommendations I have examined the above-named student and completed the preparticipation physical evaluation. The athlete does not present apparent clinical contraindications to practice and participate in the sport(s) as outlined above. A copy of the physical exam is on record in my office and can be made available to the school at the request of the parents. If conditions arise after the athlete has been cleared for participation, a physician may rescind the clearance until the problem is resolved and the potential consequences are completely explained to the athlete (and parents/quardians). Name of physician, advanced practice nurse (APN), physician assistant (PA) (print/type)__ Date of exam Address Phone _

©2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

Signature of physician, APN, PA

■ PREPARTICIPATION PHYSICAL EVALUATION CLEARANCE FORM

Name	Sex 🗆 M 🗆 F Age Date of birth	
☐ Cleared for all sports without restriction		
☐ Cleared for all sports without restriction with recommendations for further eva	uluation or treatment for	
□ Not cleared		
☐ Pending further evaluation		
☐ For any sports		
☐ For certain sports		
Reason		
Recommendations		
EMERGENCY INFORMATION		
Allergies		
7 Horgioo		
Other information		
Other information		
HCP OFFICE STAMP	SCHOOL PHYSICIAN:	
	Reviewed on	
	(Date)	
	Approved Not Approved	
	Signature:	
I have examined the above-named student and completed the prep clinical contraindications to practice and participate in the sport(s)		
and can be made available to the school at the request of the parer	ts. If conditions arise after the athlete has been cleared fo	r participation,
the physician may rescind the clearance until the problem is resolv (and parents/guardians).	ed and the potential consequences are completely explain	ed to the athlete
Name of physician, advanced practice nurse (APN), physician assistant (PA)	Nate	1
	Phone	
Signature of physician, APN, PA		
Completed Cardiac Assessment Professional Development Module		
Date Signature		



According to NJSIAA Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

According to the New Jersey State Interscholastic Athletic Association (NJSIAA) Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

The Sports Medical Advisory Committee, which includes representatives of NJSIAA member schools as well as experts in the field of healthcare and medicine, recommends the following:

- The pain from most sports-related injuries can be managed with non-narcotic medications such as acetaminophen, nonsteroidal anti-inflammatory medications like ibuprofen, naproxen or aspirin. Read the label carefully and always take the recommended dose, or follow your doctor's instructions. More is not necessarily better when taking an over-the-counter (OTC) pain medication, and it can lead to dangerous side effects.
- Ice therapy can be utilized appropriately as an anesthetic.
- Always discuss with your physician exactly what is being prescribed for pain and request to avoid narcotics.
- Tramadol, a non-opioid analgesic in the serotonin uptake inhibitor category, is a good choice should the previously listed options be insufficient to relieve pain.
- In extreme cases, such as severe trauma or post-surgical pain, opioid pain medication should not be prescribed for more than five days at a time;
- Parents or quardians should always control the dispensing of pain medications and keep them in a safe, non-accessible location; and
- Unused medications should be disposed of immediately upon cessation of use. Ask your pharmacist about drop-off locations or home disposal kits like Deterra or Medsaway.






STATE OF NEW JERSEY DEPARTMENT OF HEALTH

NJSIAA SPORTS MEDICAL **ADVISORY COMMITTEE**





Even With Proper Training and Prevention, Sports Injuries May Occur

There are two kinds of sports injuries. Acute injuries happen suddenly, such as a sprained ankle or strained back. Chronic injuries may happen after someone plays a sport or exercises over a long period of time, even when applying overuse-preventative techniques.5

Athletes should be encouraged to speak up about injuries, coaches should be supported in injury-prevention decisions, and parents and young athletes are encouraged to become better educated about sports safety.6

What Are Some Ways to Reduce the Risk of Injury?

Half of all sports medicine injuries in children and teens are from overuse. An overuse injury is damage to a bone, muscle, ligament, or tendon caused by repetitive stress without allowing time for the body to heal. Children and teens are at increased risk for overuse injuries because growing bones are less resilient to stress. Also, young athletes may not know that certain symptoms are signs of overuse.

The best way to deal with sports injuries is to keep them from happening in the first place. Here are some recommendations to consider:



PREPARE Obtain the preparticipation physical evaluation prior to participation on a school-sponsored interscholastic or intramural athletic team or squad.



CONDITIONING Maintain a good fitness level during the season and offseason. Also important are proper warm-up and cooldown exercises.



PLAY SMART Try a variety of sports and consider specializing in one sport before late adolescence to help avoid overuse injuries.



ADEQUATE HYDRATION Keep the body hydrated to help the heart more easily pump blood to muscles, which helps muscles work efficiently.



TRAINING Increase weekly training time, mileage or repetitions no more than 10 percent per week. For example, if running 10 miles one week, increase to 11 miles the following week. Athletes should also cross-train and perform sport-specific drills in different ways, such as running in a swimming pool instead of only running on the road.



REST UP Take at least one day off per week from organized activity to recover physically and mentally. Athletes should take a combined three months off per year from a specific sport (may be divided throughout the year in one-month increments). Athletes may remain physically active during rest periods through alternative low-stress activities such as stretching, yoga or walking.



PROPER EQUIPMENT Wear appropriate and properly fitted protective equipment such as pads (neck, shoulder, elbow, chest, knee, and shin), helmets, mouthpieces, face guards, protective cups, and eyewear. Do not assume that protective gear will prevent all injuries while performing more dangerous or risky activities.

Resources for Parents and Students on Preventing Substance Misuse and Abuse

The following list provides some examples of resources:

National Council on Alcoholism and Drug Dependence - NJ promotes addiction treatment and recovery.

New Jersey Department of Human Services, Division of Mental Health and Addiction Services has a mission to decrease the abuse of alcohol, tobacco and other drugs by supporting the development of a comprehensive network of prevention, intervention and treatment services in New Jersey.

New Jersey Prevention Network includes a parent's quiz on the effects of opioids.

Operation Prevention Parent Toolkit is designed to help parents learn more about the opioid epidemic, recognize warning signs, and open lines of communication with their children and those in the community.

Parent to Parent NJ is a grassroots coalition for families and children struggling with alcohol and drug addiction.

Partnership for a Drug Free New Jersey is New Jersey's anti-drug alliance created to localize and strengthen drug-prevention media efforts to prevent unlawful drug use, especially among young people.

ReachNJ provides information for parents and families, including addiction and treatment stories.

The Science of Addiction: The Stories of Teens shares common misconceptions about opioids through the voices of teens.

Youth IMPACTing NJ is made up of youth representatives from coalitions across the state of New Jersey who have been impacting their communities and peers by spreading the word about the dangers of underage drinking, marijuana use, and other substance misuse.

- **References** ¹ Massachusetts Technical Assistance Partnership for Prevention
 - ² Centers for Disease Control and Prevention
 - ³ New Jersey State Interscholastic Athletic
- Association (NJSIAA) Sports Medical Advisory Committee (SMAC)
- ⁴ Athletic Management, David Csillan, athletic trainer, Ewing High School, NJSIAA SMAC
- ⁵ National Institute of Arthritis and Musculoskeletal and Skin Diseases
- 6 USA TODAY
- ⁷ American Academy of Pediatrics

An online version of this fact sheet developed in January 2018 is available on the New Jersey Department of Education's Alcohol, Tobacco, and Other Drug Use webpage.



Voorhees Township Public Schools Opioid Use and Misuse Educational Fact Sheet Sign Off

In accordance with *N.J.S.A.* 18A:40-41.10, public school districts, approved private schools for students with disabilities, and nonpublic schools participating in an interscholastic sports program must distribute this *Opioid Use and Misuse Educational Fact Sheet* to all student-athletes and cheerleaders. In addition, schools and districts must obtain a signed acknowledgement of receipt of the fact sheet from each student-athlete and cheerleader, and for students under age 18, the parent or guardian must also sign.

This sign-off sheet is due to the appropriate school personnel as determined by your district prior to the first official practice session of the spring 2018 athletic season (March 2, 2018, as determined by the New Jersey State Interscholastic Athletic Association) and annually thereafter prior to the student-athlete's or cheerleader's first official practice of the school year.

year.
Name of School: VOORHEES MIDDLE SCHOOL
Name of School District (if applicable): VOORHEES TOWNSHIP PUBLIC SCHOOLS
I/We acknowledge that we received and reviewed the Educational Fact Sheet on the Use and Misuse of Opioid Drugs.
Student Signature:
Parent/Guardian Signature (also needed if student is under age 18):
Date:

This form MUST be returned to the school nurse PRIOR to the first official practice ¹Does not include athletic clubs or intramural events.